

# Module 3 Divide and Conquer

Some slides borrowed from CS 240.

Thanks to Anna Lubiw and other previous CS 341 instructors.

- Divide and Conquer Algorithm Basics
- Examples from previous courses
- Recurrence Relations
- Solving Recurrences by Recursion Tree
- Solving Recurrences with the Substitution Method (“Lucky Guess”)
- More Examples

# Divide and Conquer Algorithm Basics

In previous courses, we covered a few **Divide and Conquer** algorithms but maybe didn't specifically define the term.

Divide and Conquer algorithms are broken into 3 basic steps:

- 1 **Divide** - break the problem into smaller instances of the problem
- 2 **Recurse** - use recursion to solve the smaller problems
- 3 **Conquer** - combine the results of the smaller problems to solve the initial larger problem

## Examples from previous courses

**Binary Search:** Search for an element  $k$  in a sorted array  $A$ .

Note: we may have implemented this iteratively but it has a natural recursive implementation as well.

- Compute the middle index  $m$  of  $A$  and compare with  $k$ .
- If  $k = A[m]$  then return FOUND  
Else If  $k$  is smaller than  $A[m]$  then recurse on left half of  $A$   
Else ( $k$  is larger than  $A[m]$ ) so recurse on right half of  $A$

Binary Search only recurses on one of the subproblems and simply returns what the subproblem returns.

**Analysis (worst-case):**  $T(n) = 1 + T(n/2)$  but this assumes  $n$  always divides evenly.

Rigorously,  $T(n) = 1 + \max\{T(\lfloor n/2 \rfloor), T(\lceil n/2 \rceil)\}$ .

These resolve to:  $T(n) \in O(\log n)$ .

## Examples from previous courses - Sorting

### QuickSort( $A$ ):

- Partition array based on a given pivot  $\Rightarrow O(n)$  work to divide.
- 2 Subproblems: Pivot divides  $A$  into Left and right subarrays, recurse on both.
- Conquer step is easy - does nothing if algorithm is "in-place".

### Analysis:

Worst-case:  $T(n) = O(n) + T(n - 1) \in O(n^2)$

Best-case:  $T(n) = O(n) + 2T(n/2) \in O(n \log n)$

Average Case:  $O(n \log n)$

Randomized (each pivot choice is equally likely): Expected  $O(n \log n)$

# Examples from previous courses - Sorting

## MergeSort(A):

- Divide array into left and right halves.
- 2 Subproblems: Left and right half subarrays, recurse on both.
- Merge the two sorted arrays  $\Rightarrow O(n)$  work to conquer.

## Analysis:

Best-case/Worst-case:  $T(n) = 2T(n/2) + O(n) \in O(n \log n)$

Rigorously,  $T(n) = T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + cn$

# Solving Recurrence Relations


CS 240 typically only expected you to simply expand a few steps and find the pattern but introduced the ideas of a recursion tree and the substitution method.

# Solving Recurrence Relations

CS 240 typically only expected you to simply expand a few steps and find the pattern but introduced the ideas of a recursion tree and the substitution method.

## Expected recursion-depth for QuickSort

**Goal:** Analyze expected height of recursion tree.



Define  $H(\pi, R) :=$  its height for instance  $\pi$  and outcomes  $R$ .


$$H^{exp}(n) = \max_{\pi} \sum_R \Pr(R) H(\pi, R).$$

# Solving Recurrence Relations

CS 240 typically only expected you to simply expand a few steps and find the pattern but introduced the ideas of a recursion tree and the substitution method.

## Expected recursion-depth for QuickSort

**Goal:** Analyze expected height of recursion tree.



Define  $H(\pi, R) :=$  its height for instance  $\pi$  and outcomes  $R$ .

$$H^{\text{exp}}(n) = \max_{\pi} \sum_R \Pr(R) H(\pi, R).$$

**Formula:**  $H^{\text{exp}}(n) \leq 1 + \frac{1}{n} \sum_{i=0}^{n-1} \max\{H^{\text{exp}}(i), H^{\text{exp}}(n-i-1)\}$

**Claim:**  $H^{\text{exp}}(n) \leq O(\log n)$ .

**Proof:** By Induction ...



# Recursion Tree for MergeSort

$T(n) = 2T(n/2) + cn$ , if  $n$  is even

$T(1) = 0$ , if counting number of comparisons

Recursion Tree where  $n$  is a power of 2:

Total work in the recursion tree sums to:  $c \cdot n \log n$

# Solving Mergesort

If we want to be precise, the math is not trivial:

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + (n - 1) & \text{if } n > 1 \\ 0 & \text{if } n = 1 \end{cases}$$

## Solving Mergesort

If we want to be precise, the math is not trivial:

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + (n - 1) & \text{if } n > 1 \\ 0 & \text{if } n = 1 \end{cases}$$

Resolves to  $T(n) = n \lceil \log n \rceil - 2^{\lceil \log n \rceil} + 1$

The recursion tree method finds the exact solution of the recurrence when  $n$  is a power of 2,  $T(n) \in O(n \log n)$ .

## Solving Mergesort

If we want to be precise, the math is not trivial:

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + (n - 1) & \text{if } n > 1 \\ 0 & \text{if } n = 1 \end{cases}$$

Resolves to  $T(n) = n \lceil \log n \rceil - 2^{\lceil \log n \rceil} + 1$

The recursion tree method finds the exact solution of the recurrence when  $n$  is a power of 2,  $T(n) \in O(n \log n)$ .

When given  $n$  that is not a power of 2, we can also argue that performing the analysis using  $n' =$  smallest power of 2 larger than  $n$  will also give a precise result since  $n' < 2 \cdot n$ , runtimes are typically increasing and we only want the growth rate.

## Solving Recurrences by Substitution

Also known as “Lucky Guess”, “Guess and Check”, “Guess and Prove”, ...

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + (n - 1) & \text{if } n > 1 \\ 0 & \text{if } n = 1 \end{cases}$$

Guess and prove by *Induction* that  $T(n) \leq c \cdot n \log n, \forall n \geq 1$ .

## Solving Recurrences by Substitution

Also known as “Lucky Guess”, “Guess and Check”, “Guess and Prove”, ...

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + (n - 1) & \text{if } n > 1 \\ 0 & \text{if } n = 1 \end{cases}$$

Guess and prove by *Induction* that  $T(n) \leq c \cdot n \log n, \forall n \geq 1$ .

**Base case:** For  $n = 1$ ,  $T(1) = 0$  and  $c \cdot n \log n = 0$ ; i.e.  $0 \leq 0$ .

## Solving Recurrences by Substitution

Also known as “Lucky Guess”, “Guess and Check”, “Guess and Prove”, ...

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + (n-1) & \text{if } n > 1 \\ 0 & \text{if } n = 1 \end{cases}$$

Guess and prove by *Induction* that  $T(n) \leq c \cdot n \log n, \forall n \geq 1$ .

**Base case:** For  $n = 1$ ,  $T(1) = 0$  and  $c \cdot n \log n = 0$ ; i.e.  $0 \leq 0$ .

**Induction Hypothesis:** Assume that  $T(k) \leq c \cdot k \log k$  for all  $k < n$  where  $k \geq 2$ .

**Induction Step:** One method to give a rigorous proof is to separate into even and odd cases. If  $n$  is even, we don't need floors and ceilings. If  $n$  is odd, ...

## Careful! Watchout!

$$T(n) = 2T(n/2) + n$$

Claim:  $T(n) \in O(n)$

Prove  $T(n) \leq cn, \forall n \geq n_0$ .

Induction Hypothesis: Assume  $T(k) \leq c \cdot k, \forall k < n, k \geq n_0$ .

$$\begin{aligned} T(n) &= 2T(n/2) + n \\ &\leq 2c(n/2) + n \text{ by the Induction Hypothesis} \\ &= (c + 1)n \end{aligned}$$



## Careful! Watchout!

$$T(n) = 2T(n/2) + n$$

Claim:  $T(n) \in O(n)$

Prove  $T(n) \leq cn, \forall n \geq n_0$ .

Induction Hypothesis: Assume  $T(k) \leq c \cdot k, \forall k < n, k \geq n_0$ .

$$\begin{aligned} T(n) &= 2T(n/2) + n \\ &\leq 2c(n/2) + n \text{ by the Induction Hypothesis} \\ &= (c + 1)n \end{aligned}$$

Conclusion:  $T(n) \in O(n)$ .

## Careful! Watchout!

$$T(n) = 2T(n/2) + n$$

Claim:  $T(n) \in O(n)$

Prove  $T(n) \leq cn, \forall n \geq n_0$ .

Induction Hypothesis: Assume  $T(k) \leq c \cdot k, \forall k < n, k \geq n_0$ .

$$\begin{aligned} T(n) &= 2T(n/2) + n \\ &\leq 2c(n/2) + n \text{ by the Induction Hypothesis} \\ &= (c + 1)n \end{aligned}$$

Conclusion:  $T(n) \in O(n)$ .

*The conclusion is clearly incorrect!*

The problem is the constant is continuously growing.

## Substitution - Changing the Guess

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + 1 & \text{if } n > 1 \\ 1 & \text{if } n = 1 \end{cases}$$

Guess  $T(n) \in O(n)$ .

Prove by induction that  $T(n) \leq cn$  for some constant  $c$ .

## Substitution - Changing the Guess

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + 1 & \text{if } n > 1 \\ 1 & \text{if } n = 1 \end{cases}$$

Guess  $T(n) \in O(n)$ .

Prove by induction that  $T(n) \leq cn$  for some constant  $c$ .

Induction Step:

$$\begin{aligned} T(n) &= T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + 1 \\ &\leq c \cdot \lceil \frac{n}{2} \rceil + c \cdot \lfloor \frac{n}{2} \rfloor + 1 \\ &= cn + 1 \end{aligned}$$

## Substitution - Changing the Guess

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + 1 & \text{if } n > 1 \\ 1 & \text{if } n = 1 \end{cases}$$

Guess  $T(n) \in O(n)$ .

Prove by induction that  $T(n) \leq cn$  for some constant  $c$ .

Induction Step:

$$\begin{aligned} T(n) &= T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + 1 \\ &\leq c \cdot \lceil \frac{n}{2} \rceil + c \cdot \lfloor \frac{n}{2} \rfloor + 1 \\ &= cn + 1 \end{aligned}$$

What went wrong?

## Substitution - Changing the Guess

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + 1 & \text{if } n > 1 \\ 1 & \text{if } n = 1 \end{cases}$$

Guess  $T(n) \in O(n)$ .

Prove by induction that  $T(n) \leq cn$  for some constant  $c$ .

Induction Step:

$$\begin{aligned} T(n) &= T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + 1 \\ &\leq c \cdot \lceil \frac{n}{2} \rceil + c \cdot \lfloor \frac{n}{2} \rfloor + 1 \\ &= cn + 1 \end{aligned}$$

What went wrong?

Fix? Add a lower order term: Try  $T(n) \leq cn - 1$ .

## Substitution - Changing the Guess

$$T(n) = \begin{cases} T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + 1 & \text{if } n > 1 \\ 1 & \text{if } n = 1 \end{cases}$$

Guess  $T(n) \in O(n)$ .

Prove by induction that  $T(n) \leq cn$  for some constant  $c$ .

Induction Step:

$$\begin{aligned} T(n) &= T(\lceil \frac{n}{2} \rceil) + T(\lfloor \frac{n}{2} \rfloor) + 1 \\ &\leq c \cdot \lceil \frac{n}{2} \rceil + c \cdot \lfloor \frac{n}{2} \rfloor + 1 \\ &= cn + 1 \end{aligned}$$

What went wrong?

Fix? Add a lower order term: Try  $T(n) \leq cn - 1$ .

$$T(n) \leq c \cdot \lceil \frac{n}{2} \rceil - 1 + c \cdot \lfloor \frac{n}{2} \rfloor - 1 + 1 = cn - 1$$

## Substitution - Changing Variables

$$T(n) = 2T(\lfloor \sqrt{n} \rfloor) + \log n$$

Let  $m = \log n$  so  $n = 2^m$ .

Our new recurrence relation is:

$$T(2^m) = 2T(2^{m/2}) + m$$



## Substituion - Changing Variables

$$T(n) = 2T(\lfloor \sqrt{n} \rfloor) + \log n$$

Let  $m = \log n$  so  $n = 2^m$ .

Our new recurrence relation is:

$$T(2^m) = 2T(2^{m/2}) + m$$

Let  $S(m) = T(2^m)$  so  $S(m/2) = T(2^{m/2})$ .

Then  $S(m) = 2S(m/2) + m$  which is a recurrence we know.

## Substitution - Changing Variables

$$T(n) = 2T(\lfloor \sqrt{n} \rfloor) + \log n$$

Let  $m = \log n$  so  $n = 2^m$ .

Our new recurrence relation is:

$$T(2^m) = 2T(2^{m/2}) + m$$

Let  $S(m) = T(2^m)$  so  $S(m/2) = T(2^{m/2})$ .

Then  $S(m) = 2S(m/2) + m$  which is a recurrence we know.

$S(m) \in O(m \log m)$  so  $T(2^m) \in O(m \log m)$  and

$T(n) \in O((\log n)(\log \log n))$

## Example

Some websites attempt to make suggestions (or target advertising) to you based on matching you with previous users and observing what they were interested in.

Similarity of users is based on similarity of preferences. Given a set of items  $A, B, C, D$ , a user ranks them by most favourite to least favourite.

Given 2 rankings, how do you compare similarity?

Ranking 1: B D C A

Ranking 2: A D B C

## Example

Some websites attempt to make suggestions (or target advertising) to you based on matching you with previous users and observing what they were interested in.

Similarity of users is based on similarity of preferences. Given a set of items  $A, B, C, D$ , a user ranks them by most favourite to least favourite.

Given 2 rankings, how do you compare similarity?

Ranking 1: B D C A

Ranking 2: A D B C

How many exact matches are there?

## Example

Some websites attempt to make suggestions (or target advertising) to you based on matching you with previous users and observing what they were interested in.

Similarity of users is based on similarity of preferences. Given a set of items  $A, B, C, D$ , a user ranks them by most favourite to least favourite.

Given 2 rankings, how do you compare similarity?

Ranking 1: B D C A

Ranking 2: A D B C

How many exact matches are there? 1 (only D at the same index)

Maybe try ordering of pairs? How many pairs of distinct items?

## Example

Some websites attempt to make suggestions (or target advertising) to you based on matching you with previous users and observing what they were interested in.

Similarity of users is based on similarity of preferences. Given a set of items  $A, B, C, D$ , a user ranks them by most favourite to least favourite.

Given 2 rankings, how do you compare similarity?

Ranking 1: B D C A

Ranking 2: A D B C

How many exact matches are there? 1 (only D at the same index)

Maybe try ordering of pairs? How many pairs of distinct items?  $\binom{4}{2} = 6$

Are these 2 rankings similar? How many pairs  $(X, Y)$  have the same order?

2: Both prefer B over C and D over C  $\rightarrow$  pairs BC and CD.

Not very similar - 4 pairs are inverted: AB AC AD BD, *not very similar*.

## Example: Counting Inversions

### Problem

#### *Counting Inversions*

**Instance:** Given 2 rankings of items  $\{a_1, \dots, a_n\}$

**Find:** The number of inverted pairs of items between the rankings; i.e. pairs where one ranking prefers  $a_i$  over  $a_j$  but the other prefers  $a_j$  over  $a_i$ .

Observe: If we draw edges between the same items in both rankings, the number of edge crossings is the number of inversions.

## Example: Counting Inversions

### Problem

#### *Counting Inversions*

**Instance:** Given 2 rankings of items  $\{a_1, \dots, a_n\}$

**Find:** The number of inverted pairs of items between the rankings; i.e. pairs where one ranking prefers  $a_i$  over  $a_j$  but the other prefers  $a_j$  over  $a_i$ .

Observe: If we draw edges between the same items in both rankings, the number of edge crossings is the number of inversions.

An equivalent formulation is to assign numbers to each item, then compare the order of numbers. For simplicity, assign the numbers in order to the first ranking:

B D C A  $\Rightarrow$  1 2 3 4

Using the same mapping, the second ranking becomes:

A D B C



## Example: Counting Inversions

### Problem

#### *Counting Inversions*

**Instance:** Given 2 rankings of items  $\{a_1, \dots, a_n\}$

**Find:** The number of inverted pairs of items between the rankings; i.e. pairs where one ranking prefers  $a_i$  over  $a_j$  but the other prefers  $a_j$  over  $a_i$ .

Observe: If we draw edges between the same items in both rankings, the number of edge crossings is the number of inversions.

An equivalent formulation is to assign numbers to each item, then compare the order of numbers. For simplicity, assign the numbers in order to the first ranking:

B D C A  $\Rightarrow$  1 2 3 4

Using the same mapping, the second ranking becomes:

A D B C  $\Rightarrow$  4 2 1 3

The problem of counting the number of inversions now becomes ...

## Example: Counting Inversions

The problem of counting the number of inversions now becomes:  
*count the number of pairs that are out of order in the 2nd list.*

**Brute Force:**

## Example: Counting Inversions

The problem of counting the number of inversions now becomes:  
*count the number of pairs that are out of order in the 2nd list.*

**Brute Force:** Check all  $\binom{n}{2}$  pairs, requires  $O(n^2)$  time.  
Does sorting help? Not really since we check all pairs.

**Divide and Conquer Method:** Given a list  $L = a_1, \dots, a_n$  of numbers, count the number of inversions.

## Example: Counting Inversions

The problem of counting the number of inversions now becomes:  
*count the number of pairs that are out of order in the 2nd list.*

**Brute Force:** Check all  $\binom{n}{2}$  pairs, requires  $O(n^2)$  time.  
Does sorting help? Not really since we check all pairs.

**Divide and Conquer Method:** Given a list  $L = a_1, \dots, a_n$  of numbers, count the number of inversions.

- Divide  $L$  into 2 lists at  $m = \lceil \frac{n}{2} \rceil$ :  $A = a_1, \dots, a_m$  and  $B = a_{m+1}, \dots, a_n$
- Recursively count number of inversions in  $A$  and  $B \Rightarrow$  return counts  $r_A$  and  $r_B$
- Combine the results:  $r_A + r_B + r$ . What is  $r$ ?

## Example: Counting Inversions

The problem of counting the number of inversions now becomes:  
*count the number of pairs that are out of order in the 2nd list.*

**Brute Force:** Check all  $\binom{n}{2}$  pairs, requires  $O(n^2)$  time.  
Does sorting help? Not really since we check all pairs.

**Divide and Conquer Method:** Given a list  $L = a_1, \dots, a_n$  of numbers, count the number of inversions.

- Divide  $L$  into 2 lists at  $m = \lceil \frac{n}{2} \rceil$ :  $A = a_1, \dots, a_m$  and  $B = a_{m+1}, \dots, a_n$
- Recursively count number of inversions in  $A$  and  $B \Rightarrow$  return counts  $r_A$  and  $r_B$
- Combine the results:  $r_A + r_B + r$ . What is  $r$ ?  
 $r :=$  number of inversions with one element in  $A$  and one in  $B$ ; i.e. number of pairs  $(a_i, a_j)$  with  $a_i \in A$  and  $a_j \in B$  and  $a_i > a_j$

## Example: Counting Inversions

How do we find  $r$ ?

**Count:** For each  $a_j \in B$ , count the number of items,  $r_j$ , in  $A$  that are larger than  $a_j$ ; i.e.  $r = \sum_{a_j \in B} r_j$

## Example: Counting Inversions

How do we find  $r$ ?

**Count:** For each  $a_j \in B$ , count the number of items,  $r_j$ , in  $A$  that are larger than  $a_j$ ; i.e.  $r = \sum_{a_j \in B} r_j$

Now, it would help if  $A$  and  $B$  are sorted and also, for the combine step to return a sorted list. Think about how we can modify *mergesort* to compute  $r$ ; modify the *merge* process.

When  $a_j$  is copied into the merged list (combined sorted list), determine  $r_j$  and add to  $r$ .

## Example: Counting Inversions

**Algorithm:** Sort-and-Count( $L$ ) returns a sorted  $L$  and number of inversions.

- Divide  $L$  at midpoint into  $A$  and  $B$
- Sort-and-Count( $A$ ) returns (*sorted*  $A$ ,  $r_A$ )  
Sort-and-Count( $B$ ) returns (*sorted*  $B$ ,  $r_B$ )
- $r \leftarrow 0$   
Merge( $A, B$ ) and when an element of  $B$  is chosen to merge,  
 $r \leftarrow r +$  number of elements remaining in  $A$   
return (*sorted*  $A \cup B$ ,  $r_A + r_B + r$ )

**Analysis:**



## Example: Counting Inversions

**Algorithm:** Sort-and-Count( $L$ ) returns a sorted  $L$  and number of inversions.

- Divide  $L$  at midpoint into  $A$  and  $B$
- Sort-and-Count( $A$ ) returns (*sorted*  $A$ ,  $r_A$ )  
Sort-and-Count( $B$ ) returns (*sorted*  $B$ ,  $r_B$ )
- $r \leftarrow 0$   
Merge( $A, B$ ) and when an element of  $B$  is chosen to merge,  
 $r \leftarrow r +$  number of elements remaining in  $A$   
return (*sorted*  $A \cup B$ ,  $r_A + r_B + r$ )

**Analysis:** Similar to mergesort:  $T(n) = 2T(n/2) + O(n) \in O(n \log n)$   
Better Algorithms?

## Common Recurrences

We often see recurrences of the form:

$$T(n) = aT\left(\frac{n}{b}\right) + cn^k$$

Example: Mergesort  $k = 1, a = 2, b = 2$

$$T(n) = 2T\left(\frac{n}{2}\right) + cn \in O(n \log n)$$

Example:  $k = 1, a = 1, b = 2$

$$T(n) = T\left(\frac{n}{2}\right) + cn \in O(n)$$

Example:  $k = 1, a = 4, b = 2$

$$T(n) = 4T\left(\frac{n}{2}\right) + cn \in O(n^2)$$

# Master Theorem

**Theorem:** Given

$$T(n) = aT\left(\frac{n}{b}\right) + cn^k$$

where  $a \geq 1, b > 1, c > 0, k \geq 0$ , then

$$T(n) \in \begin{cases} \Theta(n^k) & \text{if } a < b^k \text{ i.e. } \log_b a < k \\ \Theta(n^k \log n) & \text{if } a = b^k \\ \Theta(n^{\log_b a}) & \text{if } a > b^k \end{cases}$$

**Proof:** For a rigorous proof, use induction.

Less rigorous, think about the recursion tree.

# Multiplying Large Numbers

**Recall:** grade 2 method  $\Rightarrow$  multiplying two  $n$ -digit numbers  $\in O(n^2)$

How about an  $n$ -digit number with an  $m$ -digit number? *Exercise*

## **Divide and Conquer Method:**

Idea: Split numbers in half (by digits), multiply smaller components.

Easier if both have same number of digits  $\Rightarrow$  pad with 0 if necessary.

## Multiplying Large Numbers

**Recall:** grade 2 method  $\Rightarrow$  multiplying two  $n$ -digit numbers  $\in O(n^2)$

How about an  $n$ -digit number with an  $m$ -digit number? *Exercise*

### Divide and Conquer Method:

Idea: Split numbers in half (by digits), multiply smaller components.

Easier if both have same number of digits  $\Rightarrow$  pad with 0 if necessary.

**Example:** Multiply 667 (0667) with 1234

06|67  $\times$  12|34 becomes the sum of:

06  $\times$  12  $\Rightarrow$  720000 (72 shifted 4 digits)

06  $\times$  34  $\Rightarrow$  20400 (204 shifted 2 digits)

67  $\times$  12  $\Rightarrow$  80400 (804 shifted 2 digits)

67  $\times$  34  $\Rightarrow$  2278

Total sum: 823078

Use recursion until numbers are small enough: 0|6  $\times$  1|2, etc

# Multiplying Large Numbers

**Analysis:**  $T(n) = 4T(n/2) + O(n)$

Apply the Master Theorem:  $a = 4, b = 2, k = 1$  and compare  $a$  with  $b^k$

$\Rightarrow 4 > 2$ , so Case 3:  $T(n) \in \Theta(n^{\log_b a})$

# Multiplying Large Numbers

**Analysis:**  $T(n) = 4T(n/2) + O(n)$

Apply the Master Theorem:  $a = 4, b = 2, k = 1$  and compare  $a$  with  $b^k$

$\Rightarrow 4 > 2$ , so Case 3:  $T(n) \in \Theta(n^{\log_b a}) \in \Theta(n^2)$

## Karatsuba's Algorithm (1960)

**Idea:** Avoid one of the four multiplications!

Consider  $0667 \times 1234$  where  $w = 06, x = 67, y = 12, z = 34$ , then

$$wx \times yz \Rightarrow w|x \times y|z$$

$$= (10^2w + x) \times (10^2y + z)$$

$$= 10^4wy + 10^2(wz + xy) + xz$$

Don't need  $wz, xy$  individually, only the sum  $(wz + xy)$ .



## Karatsuba's Algorithm (1960)

**Idea:** Avoid one of the four multiplications!

Consider  $0667 \times 1234$  where  $w = 06, x = 67, y = 12, z = 34$ , then

$$wx \times yz \Rightarrow w|x \times y|z$$

$$= (10^2w + x) \times (10^2y + z)$$

$$= 10^4wy + 10^2(wz + xy) + xz$$

Don't need  $wz, xy$  individually, only the sum  $(wz + xy)$ .

$$(w + x) \times (y + z) = wy + (wz + xy) + xz$$

$$\Rightarrow (wz + xy) = (w + x) \times (y + z) - wy - xz$$

We already compute  $wy$  and  $xz$ . Only 3 multiplications.

# Karatsuba's Algorithm (1960)

**Idea:** Avoid one of the four multiplications!

Consider  $0667 \times 1234$  where  $w = 06, x = 67, y = 12, z = 34$ , then

$$wx \times yz \Rightarrow w|x \times y|z$$

$$= (10^2w + x) \times (10^2y + z)$$

$$= 10^4wy + 10^2(wz + xy) + xz$$

Don't need  $wz, xy$  individually, only the sum  $(wz + xy)$ .

$$(w + x) \times (y + z) = wy + (wz + xy) + xz$$

$$\Rightarrow (wz + xy) = (w + x) \times (y + z) - wy - xz$$

We already compute  $wy$  and  $xz$ . Only 3 multiplications.

## Algorithm

$$p \Rightarrow wy$$

$$q \Rightarrow xz$$

$$r \Rightarrow (w + x) \times (y + z)$$

$$\text{return } 10^4p + 10^2(r - p - q) + q$$

# Karatsuba's Algorithm (1960)

Note: Additions are only linear in number of digits,  $O(n)$

**Analysis:**  $T(n) = 3T(n/2) + O(n)$

Master Theorem:  $a = 3$ ,  $b = 2$ ,  $k = 1$  and compare  $a$  with  $b^k$

$a = 3 > b^k = 2$ , so Case 3:  $T(n) \in \Theta(n^{\log_b a})$

# Karatsuba's Algorithm (1960)

Note: Additions are only linear in number of digits,  $O(n)$

**Analysis:**  $T(n) = 3T(n/2) + O(n)$

Master Theorem:  $a = 3$ ,  $b = 2$ ,  $k = 1$  and compare  $a$  with  $b^k$

$a = 3 > b^k = 2$ , so Case 3:  $T(n) \in \Theta(n^{\log_b a})$

$T(n) \in \Theta(n^{\log_b a}) = \Theta(n^{\log_2 3}) \approx \Theta(n^{1.585})$

Better Algorithms? Asymptotically faster methods for larger  $n$ .

Schönhage and Strassen (1971):  $O(n(\log n)(\log \log n))$  (often used)

More recent:  $O(n \log n)$

# Karatsuba's Algorithm (1960)

## Implementation Concerns

- 1 Numbers of different lengths - sometimes large differences.

Example:  $A$  has  $n$  digits,  $B$  has  $m$  digits and  $n \gg m$ .

Break  $A$  into  $O(n/m)$  blocks of  $m$  digits, e.g.

$$A = 342|3794|3749|4379|4297|7294|9742 \times 3422 = B$$

# Karatsuba's Algorithm (1960)

## Implementation Concerns

- 1 Numbers of different lengths - sometimes large differences.

Example:  $A$  has  $n$  digits,  $B$  has  $m$  digits and  $n \gg m$ .

Break  $A$  into  $O(n/m)$  blocks of  $m$  digits, e.g.

$$A = 342|3794|3749|4379|4297|7294|9742 \times 3422 = B$$

Multiply each block by  $B$

Sum all products (remember to do the shifts)

Analysis:  $O((n/m)m^{\log_2 3})$  or  $O(nm^{0.585})$

# Karatsuba's Algorithm (1960)

## Implementation Concerns

- 1 Numbers of different lengths - sometimes large differences.

Example:  $A$  has  $n$  digits,  $B$  has  $m$  digits and  $n \gg m$ .

Break  $A$  into  $O(n/m)$  blocks of  $m$  digits, e.g.

$$A = 342|3794|3749|4379|4297|7294|9742 \times 3422 = B$$

Multiply each block by  $B$

Sum all products (remember to do the shifts)

Analysis:  $O((n/m)m^{\log_2 3})$  or  $O(nm^{0.585})$

- 2 Which base to use? Base 10 nice for humans. In practice, for computers, Base  $2^{64}$ .

Store large numbers as an array of 64-bit integers (unsigned long):

$$A = a_0 + a_1(2^{26}) + a_2(2^{26})^2 + \dots + a_{n-1}(2^{26})^{n-1}$$

$$\Rightarrow A = a_0|a_1|a_2|\dots|a_{n-1}$$

# Multiplying Matrices

## Problem

### *Matrix Multiplication*

**Instance:** Two  $n$  by  $n$  matrices,  $A$  and  $B$ .

**Question:** Compute the  $n$  by  $n$  matrix product  $C = AB$ .

The naive algorithm (row by column for each of  $n^2$  locations) has complexity  $\Theta(n^3)$ .

**Divide and Conquer:** Divide into Submatrices of size  $n/2 \times n/2$ .



## Matrix Multiplication - Simple Divide and Conquer

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \quad B = \begin{bmatrix} e & f \\ g & h \end{bmatrix}, \quad C = AB = \begin{bmatrix} r & s \\ t & u \end{bmatrix}$$

If  $A, B$  are  $n$  by  $n$  matrices, then  $a, b, \dots, h, r, s, t, u$  are  $\frac{n}{2}$  by  $\frac{n}{2}$  matrices, where

$$\begin{aligned} r &= a e + b g & s &= a f + b h \\ t &= c e + d g & u &= c f + d h \end{aligned}$$

requiring 8 multiplications of  $\frac{n}{2}$  by  $\frac{n}{2}$  matrices to compute  $C = AB$ .

**Analysis:**  $T(n) = 8T(n/2) + O(n^2)$

Master Theorem:  $a = 8, b = 2, k = 2$  compare  $a = 8 > b^k = 4$

$T(n) \in \Theta(n^{\log_b a}) = \Theta(n^3)$

## Strassen's Algorithm (1969)

Idea: Similar to multiplication, algebraic genius (or trickery)!  
⇒ 7 subproblems instead of 8 to compute  $C = AB$ !

Define

$$P_1 = a(f - h)$$

$$P_3 = (c + d)e$$

$$P_5 = (a + d)(e + h)$$

$$P_7 = (a - c)(e + f).$$

$$P_2 = (a + b)h$$

$$P_4 = d(g - e)$$

$$P_6 = (b - d)(g + h)$$

Then, compute

$$r = P_5 + P_4 - P_2 + P_6$$

$$t = P_3 + P_4$$

$$s = P_1 + P_2$$

$$u = P_5 + P_1 - P_3 - P_7$$

**Analysis:**  $T(n) = 7T(n/2) + O(n^2)$

Master Theorem:  $a = 7, b = 2, k = 2$  compare  $a = 7 > b^k = 4$

$T(n) \in \Theta(n^{\log_b a}) = \Theta(n^{\log_2 7}) \approx \Theta(n^{2.808})$

## Centrality of Matrix Multiplication

Suppose two  $n \times n$  matrices can be multiplied in  $O(n^\omega)$  where  $2 \leq \omega \leq 3$ .

Many other problems can also then be solved in  $O(n^\omega)$ :

- Solving  $Ax = b$
- Determinant of  $A$
- Inverse of  $A$ ,  $A^{-1}$

Many problems are at least as difficult as matrix multiplication.

Example: Reduction of triangular matrix inversion to matrix multiplication.

Compute the inverse of an  $n \times n$  upper triangular matrix  $T$ .

Divide and Conquer: Decompose  $T$  into blocks of size  $n/2$

**Analysis:**  $T(n) = 2T(n/2) + O(n^\omega)$

Master Theorem:  $a = 2, b = 2, k = \omega$  and  $a = 2 < b^k = 2^\omega \geq 4$

So,  $T(n) \in \Theta(n^\omega)$

Also, Reduction of matrix multiplication to triangular matrix inversion.

# Finding the Closest Pair

## Problem

### **Closest Pair**

**Instance:** A set of  $n$  distinct points in the plane.

**Find:** Two distinct points  $p, q$  such that the distance between  $p$  and  $q$ ,

$$d(p, q) = \sqrt{(p_x - q_x)^2 + (p_y - q_y)^2}$$

is minimized.

**Brute Force:** try all pairs,  $O(n^2)$

Special case: 1D (points on a line): sort and compare consecutive pairs:  
 $O(n \log n)$

# Closest Pair - Divide and Conquer

Idea:

- Divide points in half: left half  $Q$ , right half  $R$ , dividing line  $L$
- Recursively find closest pair in  $Q, R$
- Combine - must consider points with an endpoint on each side of  $L$

# Closest Pair - Divide and Conquer

Idea:

- Divide points in half: left half  $Q$ , right half  $R$ , dividing line  $L$
- Recursively find closest pair in  $Q$ ,  $R$
- Combine - must consider points with an endpoint on each side of  $L$

Note: To divide points, it helps to sort by  $x$ -coord - once only!  
Then extract the points you need in linear time (they will also be sorted).